

## LIMNEPHILIDAE SUB-KEY E


13 Distal section of trochanters of 2<sup>nd</sup> and 3<sup>rd</sup> legs with ventral row of long flexuous setae..... 14

- Distal section of trochanters of 2<sup>nd</sup> and 3<sup>rd</sup> legs with ventral row of dagger-like spines..... 16

14 Additional face setae (fs) of femurs confined to posterior faces of 3<sup>rd</sup> legs (occasionally 1 seta may be present on femurs of 2<sup>nd</sup> legs)...  
.....*Limnephilus grisaeus*

Note: Parietal bands (pb) and pleural band (plb) indistinct or absent. Sclerite of the 9<sup>th</sup> abdominal dorsum with at least 1 intermediate setae (is) between the innermost and outermost primary setae (IP-OP) which is almost as long as outermost primary setae (OP).

Case curved, of plant material, sometimes partly or entirely of sand grains. Shallow permanent or temporary (usually acidic) open pools in mainland and in islands of Baltic Sea. Common in south, absent in extreme north.



### Should be:

14 Additional face setae (fs) of femurs usually confined only to posterior faces of 3<sup>rd</sup> legs. **Posterior faces of 2<sup>nd</sup> legs exceptionally with 1 and rarely with 2 setae**  
.....*Limnephilus grisaeus*